

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/78703 A2

(51) International Patent Classification⁷: **A61K 31/00**

(74) Agents: FISHER, Carlos, A. et al.; Allergan Sales, Inc.,
2525 Dupont Drive, Irvine, CA 92612 (US).

(21) International Application Number: PCT/US01/11843

(22) International Filing Date: 11 April 2001 (11.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
(89)548,410 13 April 2000 (13.04.2000) US

(71) Applicant: ALLERGAN SALES, INC. [US/US]; 2525
Dupont Drive, Irvine, CA 92612 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(72) Inventors: CHOW, Ken; 20 Tidal Surf, Newport Coast,
CA 92657 (US). GIL, Daniel, W.; 2541 Point Del Mar,
Corona Del Mar, CA 92625 (US). FANG, Wenkui, Ken;
73 Peppermint Tree, Irvine, CA 92618 (US). GARST,
Michael, E.; 2627 Raqueta, Newport Beach, CA 92660
(US). WHEELER, Larry, A.; 18 Valley View, Irvine, CA
92612 (US).

(54) Title: METHODS AND COMPOSITIONS FOR MODULATING ALPHA ADRENERGIC RECEPTOR ACTIVITY

(57) Abstract: Methods and compositions for the treatment of pain. Particularly disclosed are new compositions for the treatment of chronic pain, and methods for their use.

WO 01/78703 A2

METHODS AND COMPOSITIONS FOR MODULATING ALPHA
ADRENERGIC RECEPTOR ACTIVITY

5

BACKGROUND OF THE INVENTION

Human adrenergic receptors are integral membrane proteins which have been classified into two broad classes, the alpha and the beta adrenergic receptors. Both types mediate the action of the peripheral sympathetic nervous system upon binding of catecholamines, norepinephrine and epinephrine.

Norepinephrine is produced by adrenergic nerve endings, while epinephrine is produced by the adrenal medulla. The binding affinity of adrenergic receptors for these compounds forms one basis of the classification: alpha receptors tend to bind norepinephrine more strongly than epinephrine and much more strongly than the synthetic compound isoproterenol. The preferred binding affinity of these hormones is reversed for the beta receptors. In many tissues, the functional responses, such as smooth muscle contraction, induced by alpha receptor activation are opposed to responses induced by beta receptor binding.

Subsequently, the functional distinction between alpha and beta receptors was further highlighted and refined by the pharmacological characterization of these receptors from various animal and tissue sources. As a result, alpha and beta adrenergic receptors were further subdivided into α_1 , α_2 , β_1 , and β_2 subtypes.

Functional differences between α_1 and α_2 receptors have been recognized, and compounds which exhibit selective binding between these two subtypes have been developed. Thus, in WO 92/0073, the selective ability of the R(+) enantiomer of terazosin to selectively bind to adrenergic receptors of the α_1 subtype was reported. The α_1/α_2 selectivity of this compound was disclosed as being significant because agonist stimulation of the α_2 receptors was said to inhibit secretion of epinephrine and norepinephrine, while antagonism of the α_2 receptor was said to increase secretion of these hormones. Thus, the use of non-selective alpha-adrenergic blockers, such as phenoxybenzamine and phentolamine, was said to be limited by their α_2 adrenergic receptor mediated induction of increased plasma catecholamine concentration and the attendant physiological sequelae (increased heart rate and smooth muscle contraction).

For a general background on the α -adrenergic receptors, the reader's attention is directed to Robert R. Ruffolo, Jr., α -Adrenoreceptors: Molecular Biology, Biochemistry and Pharmacology, (Progress in Basic and Clinical Pharmacology series, Karger, 1991), wherein the basis of α_1/α_2 subclassification, the molecular biology, signal transduction, agonist structure-activity relationships, receptor functions, and therapeutic applications for compounds exhibiting α -adrenergic receptor affinity was explored.

The cloning, sequencing and expression of alpha receptor subtypes from animal tissues has led to the subclassification of the α_1 adrenoreceptors into α_{1A} , α_{1B} , and α_{1D} . Similarly, the α_2 adrenoreceptors have
5 also been classified α_{2A} , α_{2B} , and α_{2C} receptors. Each α_2 receptor subtype appears to exhibit its own pharmacological and tissue specificities. Compounds having a degree of specificity for one or more of these subtypes may be more specific therapeutic
10 agents for a given indication than an α_2 receptor pan-agonist (such as the drug clonidine) or a pan-antagonist.

Among other indications, such as the treatment of glaucoma, hypertension, sexual dysfunction, and
15 depression, certain compounds having alpha 2 adrenergic receptor agonist activity are known analgesics. However, many compounds having such activity do not provide the activity and specificity desirable when treating disorders modulated by alpha-
20 2 adrenoreceptors. For example, many compounds found to be effective agents in the treatment of pain are frequently found to have undesirable side effects, such as causing hypotension and sedation at systemically effective doses. There is a need for new
25 drugs that provide relief from pain without causing these undesirable side effects. Additionally, there is a need for agents which display activity against pain, particularly chronic pain, such as chronic neuropathic and visceral pain.

British Patent 1 499 485, published February 1, 1978 describes certain thiocarbamide derivatives; some of these are said to be useful in the treatment of conditions such as hypertension, depression or
5 pain.

OBJECTS OF THE INVENTION

It is an object of the invention to provide
10 compounds and compositions useful in treating disorders modulated by alpha-2 adrenoreceptors.

It is an object of this invention to provide novel compounds having substantial analgesic activity in the treatment of chronic pain, regardless of
15 origin. Chronic pain may be, without limitation, visceral, inflammatory or neuropathic in origin. Such chronic pain may arise as a result of, or be attendant to, conditions including without limitation: arthritis, (including rheumatoid
20 arthritis), spondylitis, gouty arthritis, osteoarthritis, juvenile arthritis, and autoimmune diseases including, without limitation, lupus erythematosus.

These compositions can also be used within the
25 context of the treatment of chronic gastrointestinal inflammations, Crohn's disease, gastritis, irritable bowel disease (IBD) and ulcerative colitis; and in treatment of visceral pain, including pain caused by cancer or attendant to the treatment of cancer as,
30 for example, by chemotherapy or radiation therapy.

These compositions can be used within the context of the treatment of other chronic pain symptoms, and especially in the treatment of chronic forms of neuropathic pain, in particular, without
5 limitation, neuralgia, herpes, deafferentation pain, and diabetic neuropathies.

It is also an object of this invention to provide novel compounds for treating ocular disorders, such as ocular hypertension, glaucoma,
10 hyperemia, conjunctivitis and uveitis.

It is also an object of this invention to provide novel compounds for treating the pain associated with substance abuse and/or withdrawal.


It is a still further object of this invention
15 to provide such compounds which have good activity when delivered by peroral, parenteral, intranasal, ophthalmic, and/or topical dosing, or injection.

It is also an object of this invention to provide methods of treating pain through the
20 therapeutic administration of the compounds disclosed herein.

It is further an object of the present invention to provide methods of treating conditions known to be susceptible to treatment through alpha 2 adrenergic
25 receptors.

SUMMARY OF THE INVENTION

The present invention is directed to
30 compounds having the formula:



wherein R₁ is F or H, R₂ is Cl or H, and R₃ is F or H;
and wherein if R₁ is F then R₂ and R₃ are both H; and
5 if R₁ is H then R₂ is Cl and R₃ is F, and alkyl esters
thereof, and pharmaceutically acceptable salts of
these compounds.

The invention is also directed to methods of
treating pain, particularly chronic pain, through the
10 administration of pharmaceutically effective amounts
of compounds of the above structure.


Further, the invention is directed to methods of
treating glaucoma through the administration of a
pharmaceutically effective amount of these compounds.

15

DETAILED DESCRIPTION OF THE INVENTION


In one aspect, the present invention is
directed to compounds of Formula 1:

20




wherein R_1 is F or H, R_2 is Cl or H, and R_3 is F or H; and wherein if R_1 is F then R_2 and R_3 are both H; and if R_1 is H then R_2 is Cl and R_3 is F, and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

Preferred compounds corresponding to this structure are the following compound (hereinafter termed Formula 2):



Formula 2

and the following compound (hereinafter termed Formula 3):



Formula 3

and their alkyl esters, and pharmaceutically acceptable derivatives and/or salts of these compounds.

Applicants have discovered that these compounds activate α_2 receptors, particularly α_{2B} receptors. Additionally, these compounds act as a highly

effective analgesic, particularly in chronic pain models, with minimal undesirable side effects, such as sedation and cardiovascular depression, commonly seen with agonists of the α_2 receptors.

5 Such compounds may be administered at pharmaceutically effective dosages. Such dosages are normally the minimum dose necessary to achieve the desired therapeutic effect; in the treatment of chronic pain, this amount would be roughly that
10 necessary to reduce the discomfort caused by the pain to tolerable levels. Generally, such doses will be in the range 1-1000 mg/day; more preferably in the range 10 to 500 mg/day. However, the actual amount of the compound to be administered in any given case will be
15 determined by a physician taking into account the relevant circumstances, such as the severity of the pain, the age and weight of the patient, the patient's general physical condition, the cause of the pain, and the route of administration.

20 The compounds are useful in the treatment of pain in a mammal; particularly a human being. Preferably, the patient will be given the compound orally in any acceptable form, such as a tablet, liquid, capsule, powder and the like. However, other
25 routes may be desirable or necessary, particularly if the patient suffers from nausea. Such other routes may include, without exception, transdermal, parenteral, subcutaneous, intranasal, intrathecal, intramuscular, intravenous, and intrarectal modes of
30 delivery. Additionally, the formulations may be designed to delay release of the active compound over

a given period of time, or to carefully control the amount of drug released at a given time during the course of therapy.

Another aspect of the invention is drawn to
5 therapeutic compositions comprising the compounds of Formula 1 and alkyl esters and pharmaceutically acceptable derivatives and/or salts of these compounds and a pharmaceutically acceptable
excipient. Such an excipient may be a carrier or a
10 diluent; this is usually mixed with the active compound, or permitted to dilute or enclose the active compound. If a diluent, the carrier may be solid, semi-solid, or liquid material that acts as a
excipient or vehicle for the active compound. The
15 formulations may also include wetting agents, emulsifying agents, preserving agents, sweetening agents, and/or flavoring agents. If used as in an ophthalmic or infusion format, the formulation will usually contain one or more salt to influence the
20 osmotic pressure of the formulation.

In another aspect, the invention is directed to methods for the treatment of pain, particularly chronic pain, through the administration of a compound of Formula 1, and pharmaceutically
25 acceptable alkyl esters, salts, and derivatives thereof to a mammal in need thereof. As indicated above, the compound will usually be formulated in a form consistent with the desired mode of delivery.

It is known that chronic pain (such as pain from
30 cancer, arthritis, and many neuropathic injuries) and acute pain (such as that pain produced by an

immediate mechanical stimulus, such as tissue section, pinch, prick, or crush) are distinct neurological phenomena mediated to a large degree either by different nerve fibers and neuroreceptors
5 or by a rearrangement or alteration of the function of these nerves upon chronic stimulation. Sensation of acute pain is transmitted quite quickly, primarily by afferent nerve fibers termed C fibers, which normally have a high threshold for mechanical,
10 thermal, and chemical stimulation. While the mechanisms of chronic pain are not completely understood, acute tissue injury can give rise within minutes or hours after the initial stimulation to secondary symptoms, including a regional reduction in
15 the magnitude of the stimulus necessary to elicit a pain response. This phenomenon, which typically occurs in a region emanating from (but larger than) the site of the original stimulus, is termed hyperalgesia. The secondary response can give rise
20 to profoundly enhanced sensitivity to mechanical or thermal stimulus.

The A afferent fibers ($A\beta$ and $A\delta$ fibers) can be stimulated at a lower threshold than C fibers, and appear to be involved in the sensation of chronic
25 pain. For example, under normal conditions, low threshold stimulation of these fibers (such as a light brush or tickling) is not painful. However, under certain conditions such as those following nerve injury or in the herpesvirus-mediated condition
30 known as shingles the application of even such a light touch or the brush of clothing can be very

painful. This condition is termed allodynia and appears to be mediated at least in part by A β afferent nerves. C fibers may also be involved in the sensation of chronic pain, but if so it appears
5 clear that persistent firing of the neurons over time brings about some sort of change which now results in the sensation of chronic pain.

By "acute pain" is meant immediate, usually high threshold, pain brought about by injury such as a
10 cut, crush, burn, or by chemical stimulation such as that experienced upon exposure to capsaicin, the active ingredient in chili peppers.

By "chronic pain" is meant pain other than acute pain, such as, without limitation, neuropathic pain,
15 visceral pain (including that brought about by Cron's disease and irritable bowel syndrome (IBS)), and referred pain.

EXAMPLES

20

Example 1: Synthesis of 1-(3-chloro-2-fluorobenzyl)-3-(2-hydroxyethyl)-thiourea (Formula 2)

One molar equivalent of 3-chloro-2-fluoro-benzyl
25 bromide (commercially available from e.g., Lancaster Synthesis, Ltd.) is permitted to react with 2 molar equivalents of potassium isothiocyanate in dimethylformamide (DMF) containing 0.5 molar equivalent of NaI at 90° C for 5 hours with stirring
30 to yield 3-chloro-2-fluorobenzyl isothiocyanate. The reaction mixture is permitted to cool to room

temperature, and the solution is diluted with H₂O and extracted with ether. The ether phase containing the product is removed and the reaction mixture extracted twice more with fresh ether. The ether
5 phases are combined and the product is concentrated in a Speed Vac[®] vacuum centrifuge (using house vacuum) set in a water bath at about 45° C. When the ether has evaporated, the unpurified 3-chloro-2-fluorobenzyl isothiocyanate is a viscous liquid.
10 3.57 g of this compound is mixed with 3 molar equivalents of ethanolamine in acetonitrile, and a catalytic amount (less than 1%) of DMAP (N-N-dimethyl amino pyridine) is added. The reaction mixture is incubated for 14 hours at room temperature with
15 constant stirring. The resulting solution is then concentrated using the Speed Vac[®] vacuum centrifuge in a 60° C-70° C water bath.


The product, 1-(3-chloro-2-fluorobenzyl)-3-(2-hydroxyethyl)-thiourea, is purified by liquid
20 chromatography using 200-300 mesh silica gel in a glass column. The concentrated reaction solution is applied to the column and the column washed with three column volumes of Solvent A (50% ethyl acetate/50% hexanes). The product is then eluted
25 using 2-3 column volumes of Solvent B (10% methanol/90% ethyl acetate). The eluted product is again concentrated in a Speed Vac[®] vacuum centrifuge to remove the solvent. The product is then permitted to stand at room temperature, where it crystallizes

spontaneously. The crystals are stored in the freezer at -78°C.

The product has the following spectroscopic characteristics: ^1H NMR (D_6 DMSO, 300 MHz) δ 7.98 (br s, 1 H), 7.63 (br s, 1 H), 7.46 (t, $J=3.9$ Hz, 1 H), 7.32-7.18 (m, 2 H), 4.78 (br s, 1H), 4.72 (d, $J=3.9$ Hz, 2 H), 3.47 (br s, 4 H).

In order to compare the biological activity of 1-(3-chloro-2-fluorobenzyl)-3-(2-hydroxyethyl)-thiourea with that of the 2-fluorobenzyl derivative (FORMULA 4) and the 4-fluorobenzyl derivative (FORMULA 3), FORMULA 4 is synthesized using 2-fluoro-benzyl bromide (also commercially available) as the starting material. FORMULA 3 is synthesized using commercially purchased 4-fluorobenzyl isothiocyanate. Other synthetic steps are analogous to those used above to synthesize the compound of FORMULA 2.

The 2-fluorobenzyl isothiurea derivative (hereinafter termed FORMULA 4) has the following formula:



Formula 4

The physiological activity of these compounds was tested using four models: a rat locomotor model

to assess sedation, an assay of cardiovascular activity in monkeys, a rat thermal paw withdrawal assay (Dirig et al., *J. Neurosci. Methods* 76:183-191 (1997) to test the alleviation of acute pain, and the
5 rat spinal nerve ligation allodynia model (Kim and Chung, *Pain* 50:355-363 (1992) to assess the alleviation of neuropathic pain and central sensitization typical of chronic pain. As is known to those of skill in the art, these tests are
10 established pharmacological methods for determining sedation, cardiovascular effects, acute pain and chronic pain, respectively, of pharmaceutical agents.

15 Example 2: Sedative Activity

To test sedation, six male Sprague-Dawley rats were given up to 3 mg/kg of each compound in a saline or DMSO vehicle by intraperitoneal injection (i.p.).
20 Sedation was graded 30 minutes following administration of the drug by monitoring locomotor skills as follows.

The Sprague-Dawley rats are weighed and 1 ml/kg body weight of an appropriate concentration (ie. 3
25 mg/ml for a final dose of 3 mg/kg) drug solution is injected intraperitoneally. FORMULA 3 is formulated in approximately 10% DMSO and FORMULA 2 and FORMULA 4 are formulated in 50% DMSO. The results are compared to 29 historical controls that were injected with 1
30 ml/kg saline or 50% DMSO. Rat activity is then determined 30 minutes after injection of the drug

solution. Rats are placed in a dark covered chamber and a digicom analyzer (Omnitech Electronic) quantitates their exploratory behavior for a five-minute period. The machine records each time the rat interrupts an array of 32 photoelectric beams in the X and Y orientation.

The results show that, in comparison to the appropriate vehicle controls, none of the compounds caused a statistically significant reduction in the exploratory activity of the rats. FORMULA 2 and FORMULA 3 were tested at 1 mg/kg and FORMULA 4 was tested at 3 mg/kg. Thus, the compounds are not sedating.

15

Example 3: Effects on Cardiovascular System

To test the effect of the compounds on the cardiovascular system, six cynomolgus monkeys were given 500 µg/kg of each compound by intravenous injection (i.v.). The effects of each compound on the animals' blood pressure and heart rate was measured at time intervals from 30 minutes to six hours following administration of the drug. The peak change from a baseline measurement taken 30 minutes before drug administration is recorded using a blood pressure cuff modified for use on monkeys.

The monkeys are weighed (approximately 4 kg) and an appropriate volume (0.1 ml/kg) of a 5 mg/ml solution of each compound formulated in 10% DMSO is injected into the cephalic vein in the animals' arms.

Cardiovascular measurements are made with a BP 100S automated sphygmomanometer (Nippon Colin, Japan) at 0.5, 1, 2, 4 and 6 hours.

The results show that, in comparison to the predrug control, none of the compounds have any detectable effect on the cardiovascular system.

Example 4: Alleviation of Acute Pain

Models to measure sensitivity to acute pain have typically involved the acute application of thermal stimuli; such a stimulus causes a programmed escape mechanism to remove the affected area from the stimulus. The proper stimulus is thought to involve the activation of high threshold thermoreceptors and C fiber dorsal root ganglion neurons that transmit the pain signal to the spinal cord.

The escape response may be "wired" to occur solely through spinal neurons, which receive the afferent input from the stimulated nerve receptors and cause the "escape" neuromuscular response, or may be processed supraspinally - that is, at the level of the brain. A commonly used method to measure nociceptive reflexes involves quantification of the withdrawal or licking of the rodent paw following thermal excitation. See Dirig, D.M. et al., *J. Neurosci. Methods* 76:183-191 (1997) and Hargreaves, K. et al., *Pain* 32:77-88 (1988), hereby incorporated by reference herein.

In a variation of this latter model, male Sprague-Dawley rats were tested by being placed on a

commercially available thermal stimulus device constructed as described in Hargreaves et al. This device consists of a box containing a glass plate. The nociceptive stimulus is provided by a focused projection bulb that is movable, permitting the stimulus to be applied to the heel of one or both hindpaws of the test animal. A timer is actuated with the light source, and the response latency (defined as the time period between application of the stimulus and an abrupt withdrawal of the hindpaw) is registered by use of a photodiode motion sensor array that turns off the timer and light. Stimulus strength can be controlled by current regulation to the light source. Heating is automatically terminated after 20 seconds to prevent tissue damage.

Four test animals per group were weighed (approximately 0.3 kg) and injected intraperitoneally (i.p.) with 1 ml/kg of each compound formulated in approximately 50% dimethylsulfoxide (DMSO) vehicle. Animals received a 0.3 mg/kg and a 3 mg/kg dose of the three compounds. Rats were acclimated to the test chamber for about 15 minutes prior to testing. The paw withdrawal latency was measured at 30, 60 and 120 minutes after drug administration. The right and left paws were tested 1 minute apart, and the response latencies for each paw were averaged. Stimulus intensity was sufficient to provide a temperature of 45-50 degrees centigrade to each rat hindpaw.

The results show that none of the compounds provide analgesic effects in this bioassay of acute

pain. The response latencies for rats treated with the compounds were not statistically different from the response latencies of the rats treated with vehicle alone.

5

Example 5: Alleviation of Chronic Pain

A model for chronic pain (in particular peripheral neuropathy such as causalgia) involves the
10 surgical ligation of the L5 (and optionally the L6) spinal nerves on one side in experimental animals. Rats recovering from the surgery gain weight and display a level of general activity similar to that of normal rats. However, these rats develop
15 abnormalities of the foot, wherein the hindpaw is moderately everted and the toes are held together. More importantly, the hindpaw on the side affected by the surgery appears to become sensitive to pain from low-threshold mechanical stimuli, such as that
20 producing a faint sensation of touch in a human, within about 1 week following surgery. This sensitivity to normally non-painful touch is called "tactile allodynia" and lasts for at least two months. The response includes lifting the affected
25 hindpaw to escape from the stimulus, licking the paw and holding it in the air for many seconds. None of these responses is normally seen in the control group.

Rats are anesthetized before surgery. The
30 surgical site is shaved and prepared either with betadine or Novacaine. Incision is made from the

thoracic vertebra XlIII down toward the sacrum.
Muscle tissue is separated from the spinal vertebra
(left side) at the L4 - S2 levels. The L6 vertebra is
located and the transverse process is carefully
5 removed with a small rongeur to expose the L4 - L6
spinal nerves. The L5 and L6 spinal nerves are
isolated and tightly ligated with 6-0 silk thread.
The same procedure is done on the right side as a
control, except no ligation of the spinal nerves is
10 performed.

A complete hemostasis is confirmed, then the
wounds are sutured. A small amount of antibiotic
ointment is applied to the incised area, and the rat
is transferred to the recovery plastic cage under a
15 regulated heat-temperature lamp. On the day of the
experiment, at least seven days after the surgery,
six rats per test group are administered the test
drugs by intraperitoneal (i.p.) injection or oral
gavage. For i.p. injection, the compounds are
20 formulated in approximately 50% DMSO and given in a
volume of 1 ml/kg body weight. FORMULA 2 was tested
at doses ranging between 1 and 300 μ g /kg, FORMULA 3
was tested at doses between 0.1 and 3 mg/kg and
FORMULA 4 was tested at doses of 0.3 and 3 mg/kg.
25 FORMULA 2 was also administered by oral gavage at
doses of 0.1, 0.3 and 1 mg/kg body weight to 24-hour
fasted rats. A volume equal to 1 ml/kg body weight
of an appropriate concentration (ie. 1 mg/ml for a 1
mg/kg dose) of FORMULA 2 formulated in approximately
30 50% DMSO was injected using an 18-gauge, 3-inch

gavage needle that is slowly inserted through the esophagus into the stomach.

Tactile allodynia is measured prior to and 30 minutes after drug administration using von Frey hairs that are a series of fine hairs with incremental differences in stiffness. Rats are placed in a plastic cage with a wire mesh bottom and allowed to acclimate for approximately 30 minutes. The von Frey hairs are applied perpendicularly through the mesh to the mid-plantar region of the rats' hindpaw with sufficient force to cause slight buckling and held for 6-8 seconds. The applied force has been calculated to range from 0.41 to 15.1 grams. If the paw is sharply withdrawn, it is considered a positive response. A normal animal will not respond to stimuli in this range, but a surgically ligated paw will be withdrawn in response to a 1-2 gram hair. The 50% paw withdrawal threshold is determined using the method of Dixon, W.J., *Ann. Rev. Pharmacol. Toxicol.* 20:441-462 (1980). The post-drug threshold is compared to the pre-drug threshold and the percent reversal of tactile sensitivity is calculated based on a normal threshold of 15.1 grams. The results showed that FORMULA 4 had no analgesic activity at doses up to 3 mg/kg. Surprisingly, AGN 196204 and FORMULA 2 were both able to reduce the response to the tactile stimuli that indicate tactile allodynia. FORMULA 3 reversed the allodynic pain by 34% at an i.p. dose of 0.3 mg/kg, 32% at 1 mg/kg and 26% at 3 mg/kg. FORMULA 2 reversed the allodynia by 55% at an i.p. dose of 3 µg/kg, 85% at 10 µg/kg, 90%

at 30 $\mu\text{g/kg}$, 95% at 100 $\mu\text{g/kg}$ and 92% at 300 $\mu\text{g/kg}$.
The oral doses of FORMULA 2 ranging from 0.1 to 1
mg/kg alleviated the allodynic pain by 82-86%. Thus,
FORMULA 3 and FORMULA 2 are analgesic in a model of
5 chronic pain.

10 Example 6: Treatment of Allodynia with FORMULA 3

A 50 year old male in generally good physical
condition suffers from serious pain to his upper body
due caused by contact of his skin with his clothing.
15 The patient is unable to wear clothing on his upper
body without severe pain. His symptoms suggest a
diagnosis of shingles.

The patient is given a therapeutically effective
oral dose of FORMULA 3 in capsule form as needed for
20 the treatment of pain. Following two day's treatment,
the patient reports that the allodynia resulting from
shingles is markedly reduced, and that he is able to
wear clothing on his upper body with greater comfort.

25 Example 6: Treatment of Allodynia with FORMULA 2

Same facts as in Example 5, except the patient
is given a therapeutically effective oral dose of
FORMULA 2 in capsule form as needed for the treatment
30 of pain. Following two day's treatment, the patient
reports that the allodynia resulting from shingles is
markedly reduced, and that he is able to wear
clothing on his upper body with greater comfort.

Example 7: Treatment of Visceral Pain with FORMULA 3

- 5 A 43 year old female patient suffering from
colon cancer and undergoing chemotherapy experiences
severe visceral pain associated with this primary
condition. Treatment of this pain with opiates have
been ineffective to provide substantial relief.
- 10 The patient is given a therapeutic amount of
FORMULA 3 by intravenous infusion in a
pharmaceutically acceptable vehicle. The treatment
is given twice daily. After two days the patient
reports a significant alleviation in the visceral
15 pain associated with her condition.

Example 8: Treatment of Visceral Pain with FORMULA 2


- 20 Under the same facts as Example 7, except the
patient is given FORMULA 2 instead of FORMULA 3.
After two days the patient reports a significant
alleviation in the visceral pain associated with her
condition.

- 25 The examples contained herein are intended to be
exemplary only, and do not limit the scope of the
invention, which is defined by the claims that
conclude this specification.

CLAIMS


We claim:

- 5 1. A composition comprising a compound represented by the formula:




- wherein R₁ is F or H, R₂ is Cl or H, and R₃ is F or H;
and wherein if R₁ is F then R₂ and R₃ are both H; and
10 if R₁ is H then R₂ is Cl and R₃ is F, and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

2. The composition of claim 1 wherein the compound
15 is represented by the formula:




- and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.
- 20 3. The composition of claim 1 wherein the compound is represented by the formula:



and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

- 5 4. The composition of claim 1 wherein said composition comprises a pharmaceutically acceptable excipient for therapeutic delivery of said compound.


- 10 5. The composition of claim 4 wherein said compound is represented by the formula:



and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.


15

6. The composition of claim 4 wherein said compound is represented by the formula:




20 and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

7. A method of treating pain in a mammal in need thereof comprising the step: administering to said patient a therapeutically effective dose of a composition comprising a compound represented by the formula:




- wherein R₁ is F or H, R₂ is Cl or H, and R₃ is F or H; and wherein if R₁ is F then R₂ and R₃ are both H; and if R₁ is H then R₂ is Cl and R₃ is F, and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

8. The method of claim 6 wherein said compound is represented by the formula



- and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.
9. The method of claim 7 wherein said compound is administered orally.


10. The method of claim 6 wherein said compound is represented by the formula



and alkyl esters thereof, and pharmaceutically
5 acceptable salts of these compounds.


11. The method of claim 9 wherein said compound is administered orally.

10 12. A compound represented by the formula:




wherein R_1 is F or H, R_2 is Cl or H, and R_3 is F or H;
and wherein if R_1 is F then R_2 and R_3 are both H; and
if R_1 is H then R_2 is Cl and R_3 is F, and alkyl esters
15 thereof, and pharmaceutically acceptable salts of
these compounds.

13. A compound represented by the formula



and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

14. A compound represented by the formula



5

and alkyl esters thereof, and pharmaceutically acceptable salts of these compounds.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/78703 A3

- (51) International Patent Classification⁷: A61K 31/17, A61P 25/00, 19/00, 1/04
- (21) International Application Number: PCT/US01/11843
- (22) International Filing Date: 11 April 2001 (11.04.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/548,410 13 April 2000 (13.04.2000) US
- (71) Applicant: ALLERGAN SALES, INC. [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US).
- (72) Inventors: CHOW, Ken; 20 Tidal Surf, Newport Coast, CA 92657 (US). GIL, Daniel, W.; 2541 Point Del Mar, Corona Del Mar, CA 92625 (US). FANG, Wenkui, Ken; 73 Peppermint Tree, Irvine, CA 92618 (US). GARST, Michael, E.; 2627 Raqueta, Newport Beach, CA 92660 (US). WHEELER, Larry, A.; 18 Valley View, Irvine, CA 92612 (US).
- (74) Agents: FISHER, Carlos, A. et al.; Allergan Sales, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
21 March 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHODS AND COMPOSITIONS FOR MODULATING ALPHA ADRENERGIC RECEPTOR ACTIVITY

(57) Abstract: Methods and compositions for the treatment of pain. Particularly disclosed are new compositions for the treatment of chronic pain, and methods for their use.

WO 01/78703 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/11843

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/17 A61P25/00 A61P19/00 A61P1/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, BIOSIS, EMBASE, CHEM ABS Data, PHARMAPROJECTS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 411 615 A (WARNER LAMBERT CO) 6 February 1991 (1991-02-06) *cf. abstract, page 2, 1st para., page 3, line 15 bridging with page 4, line 57*	1-14
Y	GB 2 206 347 A (SANDOZ PRODUCTS LTD) 5 January 1989 (1989-01-05) *cf. abstract, page 7, 2nd and 3rd paras.*	1-14
Y	EP 0 068 590 A (PROCTER & GAMBLE) 5 January 1983 (1983-01-05) *cf. abstract, page 2, lines 3-14*	1-14
Y	EP 0 462 933 A (SANDOZ LTD ; SANDOZ AG (DE); SANDOZ AG (AT); SANDOZ PRODUCTS LTD (G) 27 December 1991 (1991-12-27) *cf. page 3, lines 1-53, page 8, line 51 to page 9, lines 1-10*	1-14
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *8* document member of the same patent family

Date of the actual completion of the international search

5 October 2001

Date of mailing of the international search report

15/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Stoltner, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/11843

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 20 29 967 A (AMERICAN CYANAMID COMPANY) 23 December 1970 (1970-12-23) *cf. page 1 to page 2, 1st para.* -----	1-14
Y	DE 26 10 865 A (EGYT GYOGYSZERVEGYEZETI GYAR) 30 September 1976 (1976-09-30) *cf. page 2, first para., generic formula on page 3, bridging with page 4, definitions given for the groups R4-R6* -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/11843

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0411615	A	06-02-1991	US 4988688 A	29-01-1991
			AT 99290 T	15-01-1994
			AU 6001990 A	07-02-1991
			CA 2022510 A1	03-02-1991
			DE 69005557 D1	10-02-1994
			DE 69005557 T2	21-04-1994
			DK 411615 T3	24-01-1994
			EP 0411615 A1	06-02-1991
			ES 2062220 T3	16-12-1994
			IE 902792 A1	27-02-1991
			JP 3074359 A	28-03-1991
			NO 903384 A	04-02-1991
			PT 94874 A	18-04-1991
			ZA 9006057 A	29-04-1992
GB 2206347	A	05-01-1989	AU 1846488 A	05-01-1989
			DE 3821317 A1	12-01-1989
			DK 356588 A	31-12-1988
			ES 2010284 A6	01-11-1989
			FI 883097 A	31-12-1988
			FR 2617477 A1	06-01-1989
			IT 1219676 B	24-05-1990
			JP 1026551 A	27-01-1989
			LU 87261 A1	08-03-1989
			NL 8801659 A	16-01-1989
			PT 87854 A ,B	01-07-1988
			SE 8802412 A	31-12-1988
			ZA 8804693 A	28-03-1990
EP 0068590	A	05-01-1983	US 4460602 A	17-07-1984
			CA 1191862 A1	13-08-1985
			DE 3261757 D1	14-02-1985
			EP 0068590 A1	05-01-1983
EP 0462933	A	27-12-1991	AT 106866 T	15-06-1994
			AU 645090 B2	06-01-1994
			AU 7843891 A	19-12-1991
			CA 2044797 A1	19-12-1991
			CS 9101850 A3	15-01-1992
			DE 69102352 D1	14-07-1994
			DE 69102352 T2	17-11-1994
			DK 462933 T3	15-08-1994
			EP 0462933 A1	27-12-1991
			ES 2054474 T3	01-08-1994
			FI 912903 A	19-12-1991
			HU 210683 B	28-06-1995
			IE 912055 A1	18-12-1991
			JP 4230257 A	19-08-1992
			JP 7053708 B	07-06-1995
			NZ 238571 A	26-08-1993
			PL 290703 A1	21-09-1992
			PT 98004 A	31-03-1992
			ZA 9104678 A	24-02-1993
DE 2029967	A	23-12-1970	BE 752103 A1	17-12-1970
			DE 2029967 A1	23-12-1970
			FR 2052978 A5	16-04-1971
			NL 7008856 A	21-12-1970

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/11843

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2029967 A		US 3562391 A	09-02-1971
DE 2610865 A	30-09-1976	HU 173459 B	28-05-1979
		AT 347476 B	27-12-1978
		AT 185776 A	15-05-1978
		BE 839502 A1	01-07-1976
		CA 1069931 A1	15-01-1980
		CH 619210 A5	15-09-1980
		CS 188132 B2	28-02-1979
		DD 125615 A5	04-05-1977
		DE 2610865 A1	30-09-1976
		DK 107876 A	15-09-1976
		EG 12505 A	31-07-1980
		FR 2303532 A1	08-10-1976
		GB 1499485 A	01-02-1978
		IL 49194 A	16-09-1980
		IN 143559 A1	24-12-1977
		NL 7602670 A	16-09-1976
		SE 7603242 A	15-10-1976
		SU 795462 A3	07-01-1981

THIS PAGE BLANK (USPTO)